

Elżbieta SKORBIŁOWICZ¹

MACROPHYTES AS INDICATORS OF HEAVY METALS BIO-ACCUMULATION IN UPPER NAREW RIVER

MAKROFITY JAKO WSKAŹNIKI BIOPRZYSWAJALNOŚCI METALI CIĘŻKICH W GÓRNEJ NARWI

Abstract: The study aimed at evaluating the cadmium, nickel, zinc, copper, cobalt, chromium and lead contents in bottom sediments, as well as roots and above ground parts of *Carex elata* and *Acorus calamus* in upper Narew River. Studies were carried out in summer 2008 at 10 measurement points localized on Narew River (Bondary, Ploski, Doktorce, Rzedziany, Uhowo, Bokiny Zlotoria, Siekierki and Tykocin). The metal concentrations were determined by means of AAS technique. Calculated bio-accumulation coefficients for roots of examined plants were following: *Carex elata* Co > Cu > Zn > Cd > Ni > Pb = Cr while for *Acorus calamus* Co > Cu > Zn > Cd > Ni > Cr > Pb. It can be supposed that household activity, including influences of a local transport and surface runoffs, were the main sources of examined metals deposited in bottom sediments and aquatic plants of upper Narew River. Studied plant material and bottom sediments were only slightly contaminated with cadmium.

Keywords: heavy metals, macrophytes, Narew River

Introduction

The problem of aquatic environment contamination with heavy metals is still serious despite of numerous activities associated with limitations of anthropogenic emission of these elements [1–4]. Evaluation of the aquatic environment quality includes monitoring of heavy metals contents in bottom sediments, as well as in flora and fauna living in water reservoirs or flows [5–10]. They may be a valuable source of information on qualitative and quantitative changes within rivers [11–13].

The aim of present study was to evaluate the cadmium, nickel, zinc, copper, cobalt, chromium and lead contents in bottom sediments, as well as roots and above ground parts of *Carex elata* and *Acorus calamus* in upper Narew River.

¹ Department of Technology in Engineering and Environmental Protection, Faculty of Civil and Environmental Engineering, Białystok University of Technology, ul. Wiejska 45A, 15–351 Białystok, Poland, phone: +48 79 799 60 01, email: e.skorbilowicz@pb.edu.pl

Material and methods

The study was performed in summer 2008 at 10 measurement points localized on Narew River (Bondary, Ploski, Doktorce, Rzedziany, Uhowo, Bokiny Zlotoria, Siekierki and Tykocin). Plant material and bottom sediments were experimental material subjected to determinations of total contents of cadmium, nickel, zinc, copper, chromium, cobalt and lead, as well as their soluble forms. Tussock sedge (*Carex elata*) and sweet flag (*Acorus calamus*) grew at all studied points were test plants. The bottom sediment was collected from the shore zone where the sedimentation of suspended material occurs. Samples were air-dried and sieved through a polyethylene screen (0.2 mm mesh). Fraction of $< 200 \mu\text{m}$ particle diameter was used for study, because it is applied in works associated with geochemical mapping [14]. At the same time, roots and above ground parts of tussock sedge and sweet flag were collected. After drying, each sample (roots and above ground parts) were ground using agate mill. Then, bottom sediments and plant material was digested in nitric acid in a hermetic microwave system CEM Mars-5. The bottom sediments samples were also subjected to extraction of metals soluble forms in cool $1 \text{ mol} \cdot \text{dm}^{-3}$ HCl solution [15]. Applied hydrochloric acid is one of the most commonly used to isolate the non-residual metal fraction from solid samples [16]. In addition, according to Sutherland [17] and Frankowski et al [16], diluted HCl is the most suitable eluent for determination of anthropogenic sources of metals in solid samples. The metals concentrations were determined by means of AAS technique (spectrophotometer Varian SpectraAA-100).

Classification of water sediments in Poland on a base of geochemical criteria was applied to evaluate the level of sediment contamination with heavy metals [18] and threshold values taking into account the detrimental effect of pollutants accumulated in sediments on aquatic organisms [19]. The results on the content of tested metals in the upright sedge (*Carex elata*) and sweet flag (*Acorus calamus*) are based on the dry weight of plants and compared to the literature data relating to a dry weight of plants. The physiological norm of metals contents for plants was quoted after data presented by [20].

Statistica 7.1 software was used for statistical analysis of test results: arithmetic mean, standard deviation, median and also Spearman correlation coefficients between content of metals in the bottom sediments and plants, were calculated.

Results and discussion

Concentrations of studied metals in collected bottom sediments samples from upper Narew River revealed variability due to the level of the sediment dispersion, as well as the localization of the measure point. It referred both to the total and labile forms of analyzed heavy metals. Sediments were characterized by following arithmetic mean and standard deviation (Table 1) values of their total contents: $1.52 \pm 0.07 \text{ mgCd} \cdot \text{kg}^{-1}$, $14.2 \pm 1.45 \text{ mgPb} \cdot \text{kg}^{-1}$, $47.5 \pm 15.36 \text{ mgZn} \cdot \text{kg}^{-1}$, $14.8 \pm 3.99 \text{ mgCr} \cdot \text{kg}^{-1}$, $11.9 \pm 0.99 \text{ mgNi} \cdot \text{kg}^{-1}$, $3.5 \pm 1.12 \text{ mgCu} \cdot \text{kg}^{-1}$ and $2.6 \pm 0.18 \text{ mgCo} \cdot \text{kg}^{-1}$ d.m. From a point of view of the environmental pollution, the soluble portion of metals is very important,

Table 1
Heavy metals contents in bottom sediments

River	Nearest locality	Cd		Pb		Zn		Cr		Ni		Cu		Co			
		A*	B**	A	B	A	B	A	B	A	B	A	B	A	B	A	B
		Heavy metals contents in bottom sediments [mg · kg ⁻¹ d.m.]															
Narew	Tykocin	1.51	0.36	14.1	4.1	59.1	24.3	25.1	4.3	12.7	1.1	5.7	1.6	2.3	0.9		
Narew	Siekierki	1.46	0.18	13.1	5.4	48.9	18.7	17.1	2.5	10.1	1.2	3.2	0.6	2.6	0.8		
Narew	Złotoria	1.54	0.21	13.9	4.1	36.4	20.7	13.9	0.6	10.6	0.9	2.9	1.2	2.9	0.6		
Narew	Rzedziany	1.53	0.74	14.3	6.4	30.1	21.7	10.8	0.8	12.3	2.1	3.7	1.3	2.5	0.7		
Narew	Bokiny	1.58	0.19	14.2	6.9	77.8	17.2	14.1	0.7	12.3	1.2	2.9	1.1	2.8	0.8		
Narew	Uchowo	1.63	0.55	18.2	7.1	61.9	18.6	14.5	0.9	13.4	2.4	5.3	2.1	2.4	1.1		
Narew	Doktorce	1.37	0.15	13.4	4.5	47.8	19.7	14.4	0.7	12.1	1.1	2.9	1.0	2.5	0.6		
Narew	Płoski	1.54	0.42	14.1	7.3	33.9	18.9	11.6	0.5	11.3	0.8	3.2	0.5	2.6	0.8		
Narew	Narew	1.46	0.93	13.4	4.6	47.3	19.1	13.2	0.6	12.4	1.1	3.3	1.1	2.4	1.1		
Narew	Bondary	1.53	1.02	13.6	5.8	31.6	18.9	13.5	0.7	12.1	1.2	2.1	0.5	2.5	0.8		
	Arithmetic mean	1.52	0.40	14.2	5.6	47.5	19.8	14.8	1.2	11.9	1.3	3.5	1.1	2.6	0.8		
	Standard deviation	0.07	0.32	1.45	1.25	15.36	2.00	3.99	1.22	0.99	0.52	1.12	0.50	0.18	0.17		
	Median	1.54	0.39	14.4	5.2	48.0	18.9	14.4	0.6	12.1	1.0	3.3	1.2	2.6	0.8		

* A – decomposition in HNO₃ and H₂O₂; ** B – extracted with 1 mol · dm⁻³ HCl solution.

because due to their mobility, metals can be desorbed from sediments to water, as well as be accumulated in benthos organisms [15, 20]. Performed study revealed following arithmetic mean and standard deviation (Table 1) values for labile forms: cadmium – $0.40 \pm 0.32 \text{ mgCd} \cdot \text{kg}^{-1}$, $5.6 \pm 1.25 \text{ mgPb} \cdot \text{kg}^{-1}$, $18.9 \pm 2.00 \text{ mgZn} \cdot \text{kg}^{-1}$, $1.2 \pm 1.22 \text{ mgCr} \cdot \text{kg}^{-1}$, $1.3 \pm 0.52 \text{ mgNi} \cdot \text{kg}^{-1}$, $1.1 \pm 0.50 \text{ mgCu} \cdot \text{kg}^{-1}$ and $0.8 \pm 0.17 \text{ mgCo} \cdot \text{kg}^{-1}$ d.m.

Result analysis indicated that the highest level of total contents and labile forms was found for zinc and lead, whereas the lowest concentrations were determined in the case of cadmium and cobalt. The ratios of labile to total forms of heavy metals concentrations in studied bottom sediments and expressed as a percentage could be lined up in the following sequence: $\text{Zn} > \text{Pb} > \text{Cu} > \text{Co} > \text{Cd} > \text{Ni} > \text{Cr}$. It illustrates the mobility of examined metals as well as their anthropogenic origin. Studies revealed relatively low share of chromium soluble form in its total content that in majority of examined samples, was from 4 to 10 %. Kabata-Pendias [21] reported that chromium is one of the least mobile trace elements in a natural environment. In not contaminated bottom sediments, it occurs in a stable form in 84 %. Zinc is one of the most mobile metals in a natural environment, which was also confirmed by numerous studies; its soluble form share in the total content of the bottom sediments ranged from 30 % to 80 %. The lead and copper labile forms in their total contents made up about 30 % to 40 % in examined sediments, while those of cobalt and cadmium from 25 % to 30 %, and nickel from 8 % to 18 %, because it readily forms quite durable chelate and compounds, as well as complex cations and anions. Analysis of study results from bottom sediments in upper Narew River and their comparison to sediment classification in Poland on a base of geochemical criteria [18, 19], it was found that copper and zinc contents in Zlotoria, Rzedziany, Doktorce, Ploski, Narew and Bondary localities were at the level of geochemical background, while other metals (lead, nickel, chromium and cobalt) should be classified to the I class as not contaminated sediments (Table 2).

Table 2

Geochemical classification of sediments from upper Narew River

Nearest locality	Classification of metal content in sediments						
	Cd	Pb	Zn	Cr	Ni	Cu	Co
Tykocin	II	I	I	I	I	Background	I
Siekierki	II	I	I	I	I	Background	I
Zlotoria	II	I	Background	I	I	Background	I
Rzedziany	II	I	Background	I	I	Background	I
Bokiny	II	I	I	I	I	Background	I
Uchowo	II	I	I	I	I	Background	I
Doktorce	II	I	Background	I	I	Background	I
Ploski	II	I	Background	I	I	Background	I
Narew	II	I	Background	I	I	Background	I
Bondary	II	I	Background	I	I	Background	I

Source: own work based on the Bojakowska and Sokołowska [18] and Bojakowska [19].

Cadmium was the exception, because a slight exceeding the I geochemical class occurred (Table 2). Such concentrations result mainly from the anthropogenic factors, due to which it penetrates along with the surface runoff from cultivated fields with wrong phosphorus fertilizing management [20], as well as along with the sewage; transport is also a significant source of cadmium contamination [18]. Koc et al [22] and Skorbilowicz [23] argue that the modification of chemical composition of water flowing within the catchment of Narew River results primarily from structure of agricultural performance of its surface. In addition, water of Narew River is enriched in contaminants from urban settlements and storm waters from both rural and urban areas. Mioduszewski [24] and Banaszuk [25] note that relatively intensive agriculture carried out in the vicinity of the valley, meadow management on part of the bogs, and processes of organic formations decay that take place in the upper Narew River cause significant changes in migration processes within the landscape.

Contents of heavy metals at aquatic plants can be one of the indicators of anthropopresure influences on natural environment. Heavy metal contents at aquatic plants reflect their bio-available forms concentrations in a natural environment as well as valuable complement of bottom sediments studies. Values of metal arithmetic mean and standard deviation in roots of *Carex elata* (Table 3) were: 1.9 ± 0.25 mgCd · kg⁻¹, 7.6 ± 1.04 mgPb · kg⁻¹, 78.2 ± 12.79 mgZn · kg⁻¹, 7.1 ± 3.66 mgCr · kg⁻¹, 8.6 ± 0.98 mgNi · kg⁻¹, 6.2 ± 1.98 mgCu · kg⁻¹ and 9.2 ± 0.69 mgCo · kg⁻¹ d.m., while in roots of *Acorus calamus*: 2.1 ± 0.07 mgCd · kg⁻¹, 7.7 ± 1.17 mgPb · kg⁻¹, 67.4 ± 15.23 mgZn · kg⁻¹, 8.9 ± 3.9 mgCr · kg⁻¹, 9.0 ± 1.67 mgNi · kg⁻¹, 5.4 ± 1.72 mgCu · kg⁻¹ and 9.0 ± 0.93 mgCo · kg⁻¹ d.m. (Table 4). Values of arithmetic mean and standard deviation for studied metals at above ground parts of test plants were slightly different: *Carex elata* (Table 3): 1.4 ± 0.28 mgCd · kg⁻¹, 6.8 ± 0.96 mgPb · kg⁻¹, 38.7 ± 5.94 mgZn · kg⁻¹, 4.8 ± 0.92 mgCr · kg⁻¹, 5.6 ± 1.31 mgNi · kg⁻¹, 3.1 ± 0.64 mgCu · kg⁻¹ and 6.3 ± 0.75 mgCo · kg⁻¹ d.m., whereas *Acorus calamus*: 1.4 ± 0.29 mgCd · kg⁻¹, 7.8 ± 1.13 mgPb · kg⁻¹, 56.8 ± 17.83 mgZn · kg⁻¹, 5.6 ± 3.33 mgCr · kg⁻¹, 4.4 ± 0.67 mgNi · kg⁻¹, 3.8 ± 0.98 mgCu · kg⁻¹ and 6.7 ± 0.85 mgCo · kg⁻¹ d.m. (Table 4).

Cadmium and lead are dangerous and toxic to most of life forms and more available to aquatic organisms. Distribution of these metals within the plant depends on a number of factors, including the form of the metal (ion, complex), presence of other metals, as well as the species and individual characteristics, and even a plant organ. The root usually has a higher content than shoot [26, 27]. In majority of analyzed cases, studies upon Cd and Pb confirmed that fact. According to Kabata-Pendias [28], the Cd contents in plants are very diverse and most often are in the range of 0.05–0.2 mg · kg⁻¹. The toxicity symptoms occur at amounts around 5–10 mg · kg⁻¹ for sensitive plants, and 10–30 mg · kg⁻¹ for resistant plants. In all samples, it was found that concentration of a given metal in the test plants amounted to over 0.2 mg · kg⁻¹.

The average value of bioaccumulation coefficient for studied plants in the case of Cd was 1.3 ± 0.29 (Table 5).

Contents of Pb found in *Carex elata* and *Acorus calamus* did not exceed 15 mg · kg⁻¹ (Table 3 and 4), while toxic content according to Kabata-Pendias and Pendias [20] is 30

Table 3
Heavy metals contents in *Carex elata*

River	Nearest locality	Cd		Pb		Zn		Cr		Ni		Cu		Co	
		P*	R**	P	R	P	R	P	R	P	R	P	R	P	R
Narew	Tykocin	1.7	2.6	6.9	7.3	45.1	94.6	4.9	7.7	5.3	8.6	3.1	4.8	5.2	9.9
Narew	Siekierki	1.3	1.8	6.3	6.1	34.6	93.7	5.2	15.9	7.1	7.3	2.4	8.8	6.7	8.6
Narew	Złotoria	1.3	2.2	7.5	8.9	39.2	83.7	3.7	7.2	7.2	9.7	3.1	6.7	7.1	9.3
Narew	Rzedziany	1.3	1.8	8.3	7.8	39.4	50.8	5.1	9.3	4.9	8.1	3.7	4.9	4.9	9.8
Narew	Bokiny	2.2	1.8	7.1	7.1	48.1	82.1	3.8	4.5	5.1	9.1	3.7	7.7	6.6	10.2
Narew	Uchowo	1.3	1.8	6.9	9.4	44.7	67.6	4.1	3.3	5.2	8.2	3.5	9.9	6.9	9.1
Narew	Doktorce	1.4	1.9	6.4	7.1	32.1	75.7	3.9	3.8	4.8	10.6	3.3	5.1	5.9	8.9
Narew	Płoski	1.3	1.9	6.3	7.5	29.5	80.7	4.9	6.9	7.7	8.9	1.6	3.9	6.7	7.9
Narew	Narew	1.4	2.0	8.1	8.3	36.1	72.1	6.7	5.2	3.8	8.7	3.4	5.1	6.9	8.6
Narew	Bondary	1.4	1.9	6.4	6.4	38.1	80.5	5.3	6.8	4.4	8.3	3.1	5.2	6.3	9.3
Arithmetic mean		1.4	1.9	6.8	7.6	38.7	78.2	4.8	7.1	5.6	8.58	3.10	6.2	6.3	9.2
Standard deviation		0.28	0.25	0.96	1.04	5.94	12.79	0.92	3.66	1.31	0.98	0.64	1.98	0.75	0.69
Median		1.4	1.9	6.9	7.3	38.6	80.7	4.8	6.9	5.5	8.2	3.3	5.2	6.7	9.1

* P – Heavy metals contents in above ground parts; ** R – Heavy metals contents in roots.

Table 4

Contents of heavy metals in *Acoris calamus*

River	Nearest locality	Cd		Pb		Zn		Cr		Ni		Cu		Co	
		P*	R**	P	R	P	R	P	R	P	R	P	R	P	R
Narew	Tykocin	1.1	2.2	6.1	6.5	39.8	58.3	5.1	7.8	4.4	8.3	3.8	3.6	6.8	8.1
Narew	Siekierki	1.9	1.9	7.9	7.9	74.2	87.6	7.2	19.2	5.5	10.3	4.8	8.1	6.9	9.2
Narew	Złotoria	1.4	2.1	8.7	8.2	58.9	65.2	3.8	11.7	4.6	12.6	3.5	8.9	6.1	7.3
Narew	Rzedziany	1.3	2.2	8.8	8.9	52.6	61.3	4.2	7.6	4.9	9.5	3.7	4.9	6.8	8.2
Narew	Bokiny	1.8	2.3	6.7	6.4	44.6	58.2	3.9	7.5	4.2	9.3	3.7	5.1	5.9	9.6
Narew	Uchowo	1.7	1.9	9.7	8.5	56.1	66.8	4.4	7.2	3.5	8.8	3.1	5.2	6.7	9.9
Narew	Doktorce	1.5	2.3	7.8	8.9	53.4	65.1	4.1	7.3	3.9	7.8	2.2	4.1	6.8	8.7
Narew	Płoski	1.2	2.4	8.1	8.3	45.8	59.1	3.9	6.9	4.2	8.8	3.5	5.1	5.9	9.2
Narew	Narew	1.1	2.3	7.7	8.2	43.9	51.2	4.4	6.2	3.6	8.4	3.6	4.9	6.4	10.4
Narew	Bondary	1.2	1.6	6.4	5.5	99.2	101.1	14.6	7.5	5.3	6.2	5.9	4.1	8.9	9.5
	Arithmetic mean	1.4	2.1	7.8	7.7	56.8	67.4	5.6	8.9	4.41	9.0	3.8	5.4	6.7	9.0
	Standard deviation	0.29	0.07	1.13	1.17	17.83	15.23	3.33	3.9	0.67	1.67	0.98	1.72	0.85	0.93
	Median	1.4	2.2	7.8	8.1	53.1	63.1	4.4	7.5	4.3	8.8	3.7	5.1	6.8	9.2

* P – Heavy metals contents in above ground parts; ** R – Heavy metals contents in roots.

Table 5
Bio-accumulation coefficient in *Carex elata* and *Acorus calamus* in Narew River

River	Nearest locality	Bio-accumulation coefficient													
		<i>Carex elata</i>						<i>Acorus calamus</i>							
		Cd	Pb	Zn	Cr	Ni	Cu	Co	Cd	Pb	Zn	Cr	Ni	Cu	Co
Narew	Tykocin	1.7	0.5	1.6	0.3	0.7	0.8	4.3	0.7	0.5	1.0	0.3	0.7	0.6	3.5
Narew	Siekierki	1.2	0.5	1.9	1.0	0.7	2.8	3.3	1.3	0.6	1.8	1.1	1.0	2.5	3.5
Narew	Złotoria	1.4	0.6	2.3	0.5	0.9	2.3	3.2	1.4	0.6	1.8	0.8	1.2	3.1	2.5
Narew	Rzedziany	1.2	0.5	1.4	0.9	0.7	1.3	3.9	1.4	0.6	2.0	0.7	0.8	1.3	3.3
Narew	Bokiny	1.1	0.5	1.1	0.3	0.7	2.9	3.6	1.5	0.5	0.8	0.5	0.8	1.8	3.4
Narew	Uchowo	1.1	0.5	1.1	0.2	0.6	0.5	3.7	1.2	0.5	1.1	0.5	0.7	1.0	3.3
Narew	Doktorce	1.4	0.2	1.6	0.3	0.9	1.8	3.5	1.7	0.7	1.4	0.5	0.6	1.4	1.8
Narew	Płoski	1.2	0.5	2.4	0.6	0.8	1.2	3.0	1.6	0.6	1.7	0.6	0.8	1.6	3.5
Narew	Narew	1.4	0.6	1.5	0.4	0.7	1.6	3.5	1.6	0.6	1.1	0.5	0.7	1.5	4.4
Narew	Bondary	1.2	0.5	2.5	0.5	0.7	2.5	3.7	1.0	0.4	3.2	0.6	0.5	1.9	3.8
	Arithmetic mean	1.3	0.5	1.7	0.5	0.7	1.8	3.5	1.3	0.5	1.6	0.6	0.8	1.7	3.3
	Median	1.4	0.5	1.6	0.5	0.7	1.7	3.6	1.4	0.6	1.6	0.5	0.7	1.5	3.5
	Standard deviation	0.29	0.057	0.52	0.24	0.09	0.82	0.36	0.29	0.09	0.70	0.22	0.19	0.70	0.71

$\text{mg} \cdot \text{kg}^{-1}$. Lead is hardly mobile and hardly available to plants in bottom sediments, which was confirmed by the bioaccumulation coefficient that averaged to 0.5 (Table 5).

Zinc is an element that is essential for the proper functioning of living organisms, and its excess can be harmful. It is one of the more active metals in the environment [27]. Results of the plant material analysis indicate that there is not the issue of water environment contamination by this element. Average zinc contents in two analyzed plants of *Carex elata* and *Acorus calamus* were in the range of 38.75 ± 5.94 to $78.2 \pm 12.79 \text{ mg} \cdot \text{kg}^{-1}$, respectively, which according to Kabata-Pendias and Pendias [20] indicates its physiological content. The most common zinc bioaccumulation coefficient is about 1, although studies have shown its higher values: 1.7 ± 0.52 and 1.6 ± 0.70 (Table 5).

Like for zinc, there was no contamination of the environment of upper Narew River with chrome. It is generally accepted that for highly sensitive plants, Cr is harmful if present in an amount of larger than $2 \text{ mg} \cdot \text{kg}^{-1}$, while medium-resistant plants tolerate content up to $< 20 \text{ mg} \cdot \text{kg}^{-1}$ [20]. Baker and Chesnin [29] claim that concentrations above $10 \text{ mgCr} \cdot \text{kg}^{-1}$ in plants is excessive. Only in four samples were found excessive chromium content (one sample of *Carex elata*, 3 samples of *Acorus calamus*). The average value of bioaccumulation coefficient for Cr oscillated between 0.5 and 0.6.

Kabata-Pendias and Pendias [20] as well as Baker and Chesnin [29] report that the nickel concentration above $10 \text{ mg} \cdot \text{kg}^{-1}$ in the plant is excessive. All test plant samples had natural Ni content. The bioaccumulation coefficient for *Carex elata* and *Acorus calamus* varied from weak accumulation typical for nickel to moderate one. The intensive accumulation expressed above 1, was reported in a single case at *Acorus calamus*.

Copper shows a high susceptibility to bioaccumulation from the aquatic environment [27]. In a plant, this element is accumulated mainly in roots [30], as confirmed in present study. Physiological standard for copper content in plants is $5\text{--}50 \text{ mg} \cdot \text{kg}^{-1}$ according to Kabata-Pendias and Pendias [20] or $2\text{--}20 \text{ mg} \cdot \text{kg}^{-1}$ in opinion of Market [31]. The copper content in analyzed plant material ranged from 1.6 to $9.9 \text{ mg} \cdot \text{kg}^{-1}$ and most often it remained at a level of $3.1\text{--}6.7 \text{ mg} \cdot \text{kg}^{-1}$. The range of bioaccumulation coefficient for copper was high ranging from 0.5 to 3.1.

The cobalt content in tested plants from the upper Narew River ranged 4.9 to $10.4 \text{ mg} \cdot \text{kg}^{-1}$ (with mean values from 6.3 ± 0.75 and $9.2 \pm 0.69 \text{ mg} \cdot \text{kg}^{-1}$). The toxic effects related to cobalt occur at its content range about $15\text{--}50 \text{ mg} \cdot \text{kg}^{-1}$. The bioaccumulation coefficients for cobalt in *Carex elata* and *Acorus calamus* were high and amounted to 3.5 and 3.3, respectively. The results of analyzes indicate the lack of contamination due to cobalt.

Studies performed upon the chemistry at plants of two Polish lowland rivers Ołobok and Pilawa showed higher metal contents than those in the present work; mean values were: $6.2\text{--}23.0 \text{ mgPb}$, $255\text{--}303 \text{ mgZn}$, $23\text{--}55 \text{ mgCr}$, $6.6\text{--}21.4 \text{ mgNi}$, $12.9\text{--}78 \text{ mgCu}$ and $4.0\text{--}23.0 \text{ mgCo} \cdot \text{kg}^{-1}$ [13].

Studied metals in the roots were accumulated in the following order: in *Carex elata* $\text{Zn} > \text{Co} > \text{Ni} > \text{Pb} > \text{Cr} > \text{Cu} > \text{Cd}$, and in *Acorus calamus* $\text{Zn} > \text{Co} > \text{Ni} > \text{Pb} > \text{Cr} > \text{Cu} > \text{Cd}$. In contrast, in the above ground parts, metals occurred in the following

sequence: in *Carex elata* $Zn > Co > Pb > Cr > Ni > Cu > Cd$, and in *Acorus calamus* $Zn > Pb > Co > Ni > Cr > Cu > Cd$. The lowest concentrations were observed for copper and cadmium in the roots and above ground parts, while the highest ones for zinc.

Calculated bioaccumulation coefficients were as follows in roots of the examined aquatic plants: for *Carex elata* $Co > Cu > Zn > Cd > Ni > Pb = Cr$, and for *Acorus calamus* $Co > Cu > Zn > Cd > Ni > Cr > Pb$. Overall, the largest bioaccumulation coefficients in studied plants were recorded for Co and Cu, whereas the lowest for Cr and Pb. It can be also concluded that upright sedge and sweet flag in most cases accumulate examined metals in similar ways.

Studied material in the form of numerical data on metal contents was statistically analyzed. The analyses performed showed significant statistical correlations within bottom sediments between total contents of Cr and Cu and soluble forms of these elements (Table 6). It was also statistically confirmed that the Co content in *Acorus calamus* roots depends on the concentration of Co in sediments (HCl extraction). Significant correlations between the content of Pb, Zn and Cr in the roots of aquatic plants and their contents in above ground parts were recorded as well (Table 6).

Table 6

Dependencies between heavy metals contents in bottom sediments vs plants

Dependencies	Spearman correlation coefficient	Significance level (p)
$Cr^a - Cr^b$	0.695	$p < 0.05$
$Cu^a - Cu^b$	0.690	$p < 0.05$
$Co^{Tk} - Co^b$	0.675	$p < 0.05$
$Pb^{Tk} - Pb^{Tn}$	0.737	$p < 0.05$
$Zn^{Tk} - Zn^{Tn}$	0.951	$p < 0.05$
$Cr^{Sk} - Cr^{Tk}$	0.711	$p < 0.05$
$Cr^{Tk} - Cr^{Sn}$	0.749	$p < 0.05$

^a Total content in bottom sediments; ^b soluble forms content in bottom sediments; ^{Tk} *Acorus calamus* roots; ^{Tn} *Acorus calamus* above ground parts; ^{Sk} *Carex elata* roots; ^{Sn} *Carex elata* above ground parts.

Conclusions

1. Studies upon bottom sediments of the upper Narew River showed low contents of studied metals with an exception of cadmium. The copper content and in most cases zinc, occurred at the level of geochemical background. Lead, chromium, nickel and cobalt ranged within the 1st geochemical class, while cadmium in the 2nd class.

2. The achieved results from nickel, zinc, copper, cobalt, chromium and lead analysis in plant material indicate the lack of aquatic environment contamination with these elements; the only exception was cadmium, which exceeded the natural level.

3. It has been proven that aquatic plants are characterized by a greater tendency for metals accumulating than bottom sediments with an exception of lead and chromium.

4. It has been shown that the metal contents in most of tested plants were higher in the roots than in above ground parts.

5. Elevated cadmium content may be caused by anthropogenic activities, due to which the element passes along with the surface runoff from agricultural fields, where bad economy related to phosphorus fertilizers is run and wastewaters from sewage treatment plants; transport is also a major source of cadmium.

Acknowledgement

The project was funded by S/WBiŚ/4/2011.

References

- [1] Calmano W, Hong J, Förster U. Binding and mobilization of heavy metals in contaminated sediments affected by pH and redox potential. *Water Sci Technol.* 1994;28:223-235.
- [2] Baptista Neto JA, Smith BJ, Mcallister JJ. Heavy metal concentrations in surface sediments in a near-shore environment, Jurujuba Sound, Southeast Brazil. *Environ Pollut.* 2000;109(1):1-9. DOI: 10.1016/S0269-7491(99)00233-X.
- [3] Cobelo-Garcia A, Prego R. Heavy metal sedimentary record in a Galician Ria (NWSpain): Background values and recent contamination. *Mar Pollut Bull.* 2003;46:1253-1262. DOI: 10.1016/S0025-326X(03)00168-1.
- [4] Budek L, Wardas M, Kijas A, Gembalska R. Zanieczyszczenie metalami ciężkimi środowiska rzeki Szrafy (rejon Krakowa) – porównanie stanu sprzed i po powodzi w 1997 roku. *Geologia.* 2004;30(2):175-189.
- [5] Szymanowska A, Samecka-Cymerman A, Kempers AJ. Heavy Metals in three Lakes in West Poland. *Ecotox Environ Safe.* 1999;43:21-29. DOI: 10.1006/eesa.1998.1747.
- [6] Samecka-Cymerman A, Kempers AJ. Concentrations of heavy metals and plant nutrients in water, sediments and aquatic macrophytes of anthropogenic lakes (former open cut brown coal mines) differing in stage of acidification. *Sci Total Environ.* 2001;281:87-98. DOI: [http://dx.doi.org/10.1016/S0048-9697\(01\)00838-5](http://dx.doi.org/10.1016/S0048-9697(01)00838-5).
- [7] El-Sikaily A, Khaled A, El Nemr A. Heavy metals monitoring using bivalves from Mediterranean Sea and Red Sea. *Environ Monit Assess.* 2004;98:41-58. DOI: 10.1023/B:EMAS.0000038178.98985.5d.
- [8] Jamnická G, Hrivnák R, Oťaheľová R, Skoršepa M, Valachovič M. Heavy metals content in aquatic plant species from some aquatic biotopes in Slovakia. Institute of Forest Ecology, Slovak Academy of Sciences. 2005;367-369.
- [9] Pajević S, Igić R, Krstić B, Vukov D, Borišev M, Nikolić N. Chemical compositions of aquatic macrophytes from the Danube – role in biomonitoring and bioremediation. Faculty of Sciences, Department of Biology and Ecology, University of Novi Sad, Novi Sad, Serbia i Czarnogóra. 2005;485-494. DOI: 10.1127/lr/18/2008/351.
- [10] Sadecka Z. Metale ciężkie w trzcinnie pospolitej (*Phragmites Australis*). *Przemysł Chemiczny.* 2008;5:557-562.
- [11] Skorbiłowicz E. Ocena zanieczyszczenia małych cieków Puszczy Knyszyńskiej. *Acta Agrophys.* 2004;1(2):311-320.
- [12] Samecka-Cymerman A, Kempers AJ. Toxic metals in aquatic plants surviving in surface water polluted by copper mining industry. *Ecotox Environ Safe.* 2004;59:64–69. DOI: 10.1016/j.ecoenv.2003.12.002.
- [13] Samecka-Cymerman A, Kempers AJ. Heavy Metals in Aquatic Macrophytes from Two Small Rivers Polluted by Urban, Agricultural and Textile Industry Sewages SW Poland. *Arch Environ Con Tox.* 2007;53:198-206. DOI:10.1007/s00244-006-0059-6.
- [14] Lis J, Pasieczna A. Atlas geochemiczny Polski w skali 1: 2 500 000., Warszawa: Państw Inst Geol; 1995:72 p.
- [15] Dembska G, Bolałek J, Aftanas B, Guz W, Wiśniewski S. Udział formy labilnej i całkowitej metali ciężkich w osadach z Portu Gdańskiego. Problemy analityczne badań osadów dennych. Sympozjum Naukowe Komitetu Chemii Analitycznej PAN Komisja Analizy Wody, Poznań; 2001:83-97.
- [16] Frankowski M, Siepak M, Ziola A, Novotný K, Vaculovič T, Siepak J. Vertical distribution of heavy metals in grain size fractions in sedimentary rocks: Mosina-Krajkowo water well field. Poland. *Environ Monit Assess.* 2009;155:493-507. DOI:10.1007/s10661-008-0476-2.

- [17] Sutherland RA. Comparison between nonresidua Al, Co, Cu, Fe, Mn, Ni, Pb and Zn released by a three-step sequential extraction procedure and a dilute hydrochloric acid leach for soil and road deposited sediment. *Appl Geochem*. 2005;17(4):353-365. DOI:10.1016/S0883-2927(01)00095-6.
- [18] Bojakowska I, Sokołowska G. Geochemiczne klasy czystości osadów wodnych. *Przeg Geolog*. 1998;46(1):49-54.
- [19] Bojakowska I. Kryteria oceny zanieczyszczenia osadów wodnych. *Przeg Geolog*. 2001;49(3):213-218.
- [20] Kabata-Pendias A, Pendias H. Biogeochemia pierwiastków śladowych. Warszawa: PWN; 1999:364 p.
- [21] Kabata-Pendias A. Biogeochemia chromu, niklu i glinu. In: Chrom, nikiel i glin w środowisku – problemy ekologiczne i metodyczn. Wrocław: Ossolineum; 1993:9-14.
- [22] Koc J, Nowicki A, Glińska K, Łachacz A. Kształtowanie się jakości wód w warunkach małej antropopresji na przykładzie zlewni strugi Ardung (Pojezierze Olsztyńskie). *Zesz Nauk Komitetu PAN „Człowiek i Środowisko”*. 2000;25:155-167.
- [23] Skorbiłowicz M. Czynniki i procesy kształtujące obieg składników mineralnych w wodach rzecznych zlewni górnej Narwi. *Rozp nauk nr 197*. Białystok: Oficyna Wydawnicza Politechniki Białostockiej; 2010:159 p.
- [24] Mioduszewski W. Analiza zasobów i potrzeb wodnych w zlewni górnej Narwi. In: Ochrona zasobów wodnych na przykładzie zlewni górnej Narwi. Mioduszewski W, editor. Falenty: Wyd IMUZ; 1996:7-19.
- [25] Banaszuk P. Identyfikacja procesów kształtujących skład chemiczny małego cieku w krajobrazie rolniczym na podstawie analizy czynnikowej. *Woda – Środowisko – Obszary Wiejskie*. 2004;4,1(10):103-116.
- [26] Cardwell AJ, Hawker DW, Greenway M. Metal accumulation in aquatic macrophytes from southeast Queensland, Australia. *Chemosphere*. 2002;48:653-663. DOI:10.1016/S0045-6535(02)00164-9.
- [27] Skorbiłowicz E. Studia nad rozmieszczeniem niektórych metali w środowisku wodnym zlewni górnej Narwi. *Rozp nauk nr 222*. Białystok: Oficyna Wydawnicza Politechniki Białostockiej; 2012:212 p.
- [28] Kabata-Pendias A. Kadm w środowisku – problemy ekologiczne i metodyczne. *Zesz Nauk Komitetu PAN „Człowiek i Środowisko”*. 2000;26:17-24.
- [29] Baker DE, Chesnin L. Chemical monitoring of soil environmental quality and Animal and human health. *Adv Agron*. 1975;27:305-374.
- [30] Jackson LJ. Paradigms of accumulation in rooted aquatic vascular plants. *Sci Total Environ*. 1998;219(2-3):223-231. DOI: [http://dx.doi.org/10.1016/S0048-9697\(98\)00231-9](http://dx.doi.org/10.1016/S0048-9697(98)00231-9).
- [31] Markert B. Presence and significance of naturally occurring chemical elements of the periodic system in the plant organism and consequences for future investigations on inorganic environmental chemistry in ecosystems. *Vegetation*. 1992;103:1-30. DOI: 10.1007/BF00033413.

MAKROFITY JAKO WSKAŹNIKI BIOPRZYSWAJALNOŚCI METALI CIĘŻKICH W GÓRNEJ NARWI

Katedra Technologii w Inżynierii i Ochronie Środowiska
Wydział Budownictwa i Inżynierii Środowiska, Politechnika Białostocka

Abstrakt: Celem przeprowadzonych badań było określenie zawartości kadmu, niklu, cynku, miedzi, kobaltu, chromu i ołowiu w osadach dennych oraz korzeniach i częściach nadziemnych roślin: *Carex elata* i *Acorus calamus* w górnej Narwi. Badania wykonano latem w 2008 r. w 10 przekrojach pomiarowych zlokalizowanych na rzece Narew (Bandary, Ploski, Doktorce, Rzędziany, Uhowo, Bokiny Złotoria, Siekierki i Tykocin). Stężenie metali oznaczano metodą ASA. Obliczone współczynniki bioakumulacji w korzeniach badanych roślin wodnych kształtowały się w następujący sposób: *Carex elata* Co > Cu > Zn > Cd > Ni > Pb = Cr i *Acorus calamus* Co > Cu > Zn > Cd > Ni > Cr > Pb. Można sądzić, że źródłem badanych metali zdeponowanych w osadach dennych i badanych roślinach wodnych w górnej Narwi jest działalność gospodarcza i bytowa człowieka, w tym wpływ lokalnej komunikacji oraz spływy powierzchniowe. Badany materiał roślinny i osady dennie rzeki Narew są w niewielkim stopniu zanieczyszczone kadmem.

Słowa kluczowe: metale ciężkie, makrofity, rzeka Narew